

PARABOLIC FINITE VOLUME ELEMENT EQUATIONS IN NONCONVEX POLYGONAL DOMAINS

P. CHATZIPANTELIDIS¹, R.D. LAZAROV², AND V. THOMÉE³

ABSTRACT. We study spatially semidiscrete and fully discrete finite volume element approximations of the heat equation with homogeneous Dirichlet boundary conditions in a plane polygonal domain with one reentrant corner. We show that, as a result of the singularity in the solution near the reentrant corner, the convergence rate is reduced from optimal second order, similarly to what was observed for the finite element method in the earlier work [5]. Optimal order convergence may be restored by mesh refinement near the corners of the domain.

1. INTRODUCTION

We shall consider the finite volume method, using continuous, piecewise linear approximating functions, for the model parabolic initial boundary value problem

$$(1.1) \quad \begin{aligned} u_t - \Delta u &= f(t) && \text{in } \Omega, && \text{with } u(\cdot, t) = 0 && \text{on } \partial\Omega, && \text{for } t > 0, \\ u(\cdot, 0) &= v && \text{in } \Omega, \end{aligned}$$

where Ω is a nonconvex polygonal domain in \mathbb{R}^2 . We assume for simplicity that exactly one interior angle ω is reentrant, i.e., such that $\omega \in (\pi, 2\pi)$, and set $\beta = \pi/\omega \in (\frac{1}{2}, 1)$.

In [4] we showed an $O(h^2)$ error bound in L_2 in the case of a convex Ω , and in [5] we discussed the error in the nonconvex case for the finite element method. In the latter case the error in L_2 is reduced from $O(h^2)$ to $O(h^{2\beta})$, as a result of the singularity which is present in the solution of (1.1) at the reentrant corner. In this paper we show the corresponding result for a finite volume method. We also discuss error estimations in H^1 and in the maximum-norm. The present work can be considered as a continuation of [4] and [5], and we refer to these papers for references to the literature.

The finite volume method relies on a local conservation property associated with the differential equation. Namely, integrating (1.1) over any region $V \subset \Omega$ and using Green's formula, we obtain

$$(1.2) \quad \int_V u_t dx - \int_{\partial V} \nabla u \cdot n ds = \int_V f dx, \quad \text{for } t > 0,$$

where n denotes the unit exterior normal vector to ∂V .

¹Partially supported by the Research Committee of the University of Crete under Research Project no. 2299.

²Partially supported by the US National Science Foundation under Grant CNS-0540136.

³Supported by the Institute of Applied and Computational Mathematics, FO.R.T.H, Heraklion, Greece.




FIGURE 1. *Left:* A union of triangles that have a common vertex z ; the dotted line shows the boundary of the corresponding control volume V_z . *Right:* A triangle K partitioned into the three subregions K_z .

There are various approximation strategies in the finite volume (control-volume) method. For comprehensive presentations and references to existing results and various applications we refer to the monographs [7, 9]. Here we shall study spatially semidiscrete approximations of (1.1) by the finite volume element method, which for brevity we will refer to as the finite volume method below. The approximate solution will be sought in the space of piecewise linear finite elements

$$S_h \equiv S_h(\Omega) = \{\chi \in \mathcal{C}(\Omega) : \chi|_K \text{ linear}, \forall K \in \mathcal{T}_h; \chi|_{\partial\Omega} = 0\},$$

where $\{\mathcal{T}_h\}_{0 < h < 1}$ is a family of regular triangulations of Ω , with h denoting the maximum diameter of the triangles of \mathcal{T}_h . In the sequel, for simplicity, we shall suppress the index Ω in the notation of functional spaces.

The semidiscrete finite volume approximation $u_h(t) \in S_h$, $t \geq 0$, will satisfy the relation (1.2) for V in a finite collection of subregions of Ω called control volumes, the number of which will be equal to the dimension of the finite element space S_h . These control volumes are constructed in the following way. Let z_K be the barycenter of $K \in \mathcal{T}_h$. We connect z_K with line segments to the midpoints of the edges of K , thus partitioning K into three quadrilaterals K_z , $z \in Z_h(K)$, where $Z_h(K)$ are the vertices of K . Then with each vertex $z \in Z_h = \cup_{K \in \mathcal{T}_h} Z_h(K)$ we associate a control volume V_z , which consists of the union of the subregions K_z , sharing the vertex z (see Figure 1). We denote the set of interior vertices of Z_h by Z_h^0 . The semidiscrete finite volume method is then to find $u_h(t) \in S_h$ for $t \geq 0$ such that

$$\int_{V_z} u_{h,t} dx - \int_{\partial V_z} \nabla u_h \cdot n ds = \int_{V_z} f dx, \quad \forall z \in Z_h^0, t > 0, \quad \text{with } u_h(0) = v_h,$$

with $v_h \in S_h$ a given approximation of v .

This problem may also be expressed in a weak form. For this purpose we introduce the finite dimensional piecewise constant space

$$Y_h = \{\eta \in L_2 : \eta|_{V_z} = \text{constant}, \forall z \in Z_h^0; \eta|_{V_z} = 0, \forall z \in \partial\Omega\}.$$

We now multiply the integral relation above by an arbitrary $\eta(z)$, $\eta \in Y_h$ and sum over all $z \in Z_h^0$ to obtain the Petrov–Galerkin formulation

$$(1.3) \quad (u_{h,t}, \eta) + a_h(u_h, \eta) = (f, \eta), \quad \forall \eta \in Y_h, t > 0, \quad \text{with } u_h(0) = v_h,$$

where $(v, w) = \int_{\Omega} vw \, dx$ and the bilinear form $a_h(\cdot, \cdot) : S_h \times Y_h \rightarrow \mathbb{R}$ is defined by

$$a_h(v, \eta) = - \sum_{z \in Z_h^0} \eta(z) \int_{\partial V_z} \nabla v \cdot n \, ds, \quad v \in S_h, \quad \eta \in Y_h.$$

Obviously, we can extend the definition of $a_h(v, \eta)$ for v in the fractional order Sobolev space H^{1+s} , $s > 1/2$, and using Green's formula we easily see that for $v \in H^2$,

$$(1.4) \quad a_h(v, \eta) = -(\Delta v, \eta), \quad \forall \eta \in Y_h.$$

The stationary elliptic problem corresponding to (1.1) is the Dirichlet problem,

$$(1.5) \quad -\Delta u = f \quad \text{in } \Omega, \quad \text{with } u = 0 \quad \text{on } \partial\Omega.$$

For this problem, the reentrant corner \mathcal{O} gives rise to a singularity in the solution with a leading term of the form $c(f)r^\beta \sin(\beta\theta)$, in polar coordinates centered at \mathcal{O} , even when f is smooth. This function is not in the space H^{1+s} for any $s \geq \beta$.

The finite volume method approximates the solution of (1.5) by $u_h \in S_h$ from

$$a_h(u_h, \eta) = (f, \eta), \quad \forall \eta \in Y_h,$$

and the error may be shown to satisfy

$$(1.6) \quad \|u_h - u\| \leq Ch^{2\beta} \|\Delta u\|_{H^{2\beta-1}}, \quad \text{where } \|\cdot\| = \|\cdot\|_{L_2} \text{ and } 1/2 < \beta < 1.$$

For the corresponding finite element method,

$$a(\underline{u}_h, \chi) = (f, \chi), \quad \forall \chi \in S_h, \quad \text{where } a(v, w) = \int_{\Omega} \nabla v \cdot \nabla w \, dx,$$

we have an error bound of the same order, which requires less regularity, namely

$$(1.7) \quad \|\underline{u}_h - u\| \leq C_s h^{2\beta} \|\Delta u\|_{H^{-1+s}}, \quad \text{for } \beta < s \leq 1.$$

As a guide to our analysis of (1.3) we use the corresponding finite element problem,

$$(1.8) \quad (\underline{u}_{h,t}, \chi) + a(\underline{u}_h, \chi) = (f, \chi), \quad \forall \chi \in S_h, \quad t > 0, \quad \text{with } \underline{u}_h(0) = v_h.$$

Here, in the error analysis, it is customary to split the error into two terms by

$$\underline{u}_h(t) - u(t) = (\underline{u}_h(t) - R_h u(t)) + (R_h u(t) - u(t)) = \underline{\varrho}(t) + \varrho(t).$$

where $R_h : H_0^1 \rightarrow S_h$ denotes the Ritz projection defined by

$$(1.9) \quad a(R_h v, \chi) = a(v, \chi), \quad \forall \chi \in S_h,$$

and H_0^1 denotes the H^1 Sobolev space with homogeneous boundary conditions. As a result of (1.7), we immediately have

$$(1.10) \quad \|\varrho(t)\| \leq C_s h^{2\beta} \|\Delta u(t)\|_{H^{-1+s}}, \quad \text{for } \beta < s \leq 1, \quad t > 0,$$

and we find that $\underline{\varrho}$ satisfies

$$(\underline{\varrho}_t, \chi) + a(\underline{\varrho}, \chi) = -(\varrho_t, \chi), \quad \forall \chi \in S_h, \quad t > 0.$$

This leads to an $O(h^{2\beta})$ bound also for $\|\underline{\varrho}\|$, and thus of the total error.

For the error analysis of the semidiscrete method (1.3) it would seem more natural to split the error using the finite volume elliptic projection $\tilde{R}_h : H^{1+s} \cap H_0^1 \rightarrow S_h$, $s > 1/2$, defined by

$$(1.11) \quad a_h(\tilde{R}_h v, \eta) = a_h(v, \eta), \quad \forall \eta \in Y_h,$$

and thus write

$$(1.12) \quad u_h(t) - u(t) = (u_h(t) - \tilde{R}_h u(t)) + (\tilde{R}_h u(t) - u(t)) = \tilde{\vartheta}(t) + \tilde{\varrho}(t).$$

The second term, $\tilde{\varrho}$ then represents the error in an elliptic problem whose exact solution is u and by (1.6) this term may be bounded by

$$\|\tilde{\varrho}(t)\| \leq Ch^{2\beta} \|\Delta u(t)\|_{H^{2\beta-1}}, \quad \text{for } t > 0.$$

For the first term in (1.12), $\tilde{\vartheta}(t) \in S_h$, we have

$$(1.13) \quad (\tilde{\vartheta}_t, \eta) + a_h(\tilde{\vartheta}, \eta) = -(\tilde{\varrho}_t, \eta), \quad \forall \eta \in Y_h, \quad t > 0.$$

and it follows easily that $\|\tilde{\vartheta}(t)\|$, and thus also the total error, are of order $O(h^{2\beta})$. However, as we shall see, these error bounds will require higher regularity assumptions and compatibility conditions on data than for the finite element method.

In an alternative analysis, proposed in [4], we may split the error using the finite element Ritz projection R_h , and thus write

$$(1.14) \quad u_h(t) - u(t) = (u_h(t) - R_h u(t)) + (R_h u(t) - u(t)) = \vartheta(t) + \varrho(t).$$

We then have estimate (1.10) for ϱ , whereas ϑ now satisfies the somewhat more complicated equation

$$(1.15) \quad (\vartheta_t, \eta) + a_h(\vartheta, \eta) = -(\varrho_t, \eta) - a_h(\varrho, \eta), \quad \forall \eta \in Y_h, \quad t > 0.$$

This equation also makes it possible to show an $O(h^{2\beta})$ bound for ϑ and thus for $u_h - u$. It turns out that the regularity requirements using this method, although still slightly higher than for the finite element method, are less stringent than what is needed by using the finite volume elliptic projection \tilde{R}_h .

Using the Ritz projection in the error splitting, i.e., (1.14), we also derive, as for the finite element method in [5], an $O(h^\beta)$ bound for the gradient of the error and an almost $O(h^\beta)$ global error estimate in maximum-norm. In maximum-norm, we also show an $O(h^{2\beta})$ error bound, away from the corners, and finally demonstrate that the almost optimal order $O(h^2)$ error bound may be restored by refining the triangulations near the corners. The regularity requirements for these error bounds, as in the L_2 norm estimate, are higher than those needed for the finite element method.

The following is an outline of the paper. In Section 2 we briefly recall from [5] some definitions of function spaces, regularity results, and error bounds for finite element approximations for elliptic and parabolic problems, that will be useful subsequently. The main section of the paper is Section 3, where error bounds in L_2 , and H^1 are shown together with three maximum-norm error estimates, mentioned above. In Section 4 we derive similar error bounds for a fully discrete scheme by discretizing also in time using the Backward Euler method.

As in [4] and [5], our error bounds will be expressed in terms of norms of data, together with compatibility conditions at $\partial\Omega$ for $t = 0$. This should be interpreted to mean that if the bounds are finite, then the exact solution will have enough regularity to secure the convergence rate stated. In the error bounds, C will denote constants which may depend on Ω and on geometrical properties of \mathcal{T}_h , but are independent of h and data. Several of the constants in our error and regularity bounds grow with t , and in order to not to have to account for their precise growth, we will assume throughout that $t \leq T$, for some positive T , without indicating the dependence of the constant on T . Also, in our analysis, sometimes norms in the spatial variable of fractional order occur, and to make for easier reading we often

replace such norms with norms of integral order in our statements. Further, we shall make particular choices for v_h in order to simplify the presentation. Note that by the stability of (1.3), other natural choices of v_h would give the same error bounds.

2. REVIEW OF THE ERROR ANALYSIS FOR FINITE ELEMENT APPROXIMATIONS

In this section we collect some material from [5] that we will need in our subsequent analysis, namely some definitions relating to fractional order Sobolev spaces, regularity results for the Dirichlet problem (1.5) and the parabolic model problem (1.1), and error bounds for the Ritz projection R_h . To be able to compare our new results for the finite volume solution of (1.1) with the corresponding error bounds for the finite element solution, we include some of the latter. For more details and references to the literature, we refer to [5].

Letting $H^{-1} = (H_0^1)^*$ denote the dual space of H_0^1 , with respect to the L_2 inner product, we define the variational solution of (1.5) for $f \in H^{-1}$ as the function $u \in H_0^1$ which satisfies

$$(2.1) \quad (\nabla u, \nabla \varphi) = (f, \varphi), \quad \forall \varphi \in H_0^1,$$

thus also defining the operator $\Delta : H_0^1 \rightarrow H^{-1}$. It is well-known that this problem has a unique solution, and that $\|\nabla u\| \leq \|f\|_{H^{-1}}$. In order to discuss further regularity results we shall need to use fractional order Sobolev spaces. Let H^m with norm $\|\cdot\|_{H^m}$ denote the standard Sobolev spaces of order m , with m integer. The space H^s , for s non integer, $s = m + \sigma$, $0 \leq \sigma \leq 1$, is defined by the real interpolation method, $H^s = [H^m, H^{m+1}]_{\sigma, 2}$.

Also let H_0^s , $0 \leq s \leq 1$, be a fractional order Sobolev space obtained by interpolation between L_2 and H_0^1 . Note that $H_0^s = H^s$ for $0 < s < 1/2$, which means H_0^s does not require any boundary condition for small s . Further, we denote $H^{-s} = (H_0^s)^*$, the dual space with respect to the L_2 inner product, for $0 \leq s \leq 1$.

For the error analysis of (1.8) and (1.3) we use the Hilbert spaces \dot{H}^s defined by the norms

$$\|v\|_{\dot{H}^s} = \left(\sum_{j=1}^{\infty} \lambda_j^s (v, \varphi_j)^2 \right)^{1/2}, \quad \text{for } s \geq -1, \quad v \in H^{-1}.$$

where $\{\varphi_j\}_{j=1}^{\infty}$ are the orthonormal eigenfunctions and $\{\lambda_j\}_{j=1}^{\infty}$ the corresponding eigenvalues of $-\Delta$ in Ω .

Since both \dot{H}^{-s} and H^{-s} are the uniquely defined interpolation space between L_2 and H^{-1} , we have $\dot{H}^{-s} = H^{-s}$ for $0 \leq s \leq 1$, and for $1 \leq s \leq 2$, \dot{H}^s consists of the functions $u \in H_0^1$ such that Δu is in the space H^{s-2} . Further it is obvious that $-\Delta$ gives an isomorphism between \dot{H}^{1+s} and \dot{H}^{-1+s} . Thus,

$$\|\Delta u\|_{H^{-1+s}} \leq C \|\Delta u\|_{\dot{H}^{-1+s}} = C \|u\|_{\dot{H}^{1+s}}, \quad \text{for } 0 \leq s \leq 1.$$

It is well-known that the nonconvex corner of Ω bounds the regularity of the solution u of (1.5). Thus $u \in H^{1+s}$ for $0 \leq s < \beta$ for f smooth enough, $\|u\|_{H^{1+s}} \leq C_s \|f\|_{H^{-1+s}}$, but $u \notin H^{1+\beta}$. A somewhat more sophisticated regularity result makes it possible to show the following error bounds in L_2 and energy norms, for the Ritz finite element projection R_h , defined by (1.9).

Lemma 2.1. *Let u be the solution of (1.5) or (2.1). Then, we have, with $C = C_s$, for $\beta < s \leq 1$,*

$$(2.2) \quad \|R_h u - u\| + h^\beta \|\nabla(R_h u - u)\| \leq Ch^{2\beta} \|\Delta u\|_{H^{-1+s}} \leq Ch^{2\beta} \|u\|_{\dot{H}^{1+s}}.$$

Further,

$$(2.3) \quad \|R_h u - u\| \leq Ch^\beta \|u\|_{H^1}.$$

In the maximum-norm $\|v\|_C = \sup_{x \in \Omega} |v(x)|$ the following almost $O(h^\beta)$ error estimate holds.

Lemma 2.2. *Let u be the solution of (1.5). If the triangulations \mathcal{T}_h are such that $h_{\min} \geq Ch^\gamma$ for some $\gamma > 0$, then, for any s, s_1 with $0 \leq s < s_1 < \beta$, we have, with $C = C_{s, s_1}$,*

$$\|R_h u - u\|_C \leq Ch^s \|u\|_{\dot{H}^{1+s_1}}.$$

We note that for quasiuniform triangulations the logarithmic stability estimate

$$\|R_h v\|_C \leq C \ell_h \|v\|_C, \quad \text{where } \ell_h = \max(\log(1/h), 1),$$

may be used to improve the maximum-norm convergence rate to $O(h^\beta \ell_h)$.

Away from the corners of the domain Ω , the convergence in maximum-norm is of the same order $O(h^{2\beta})$ as in the global L_2 error estimate. For this we quote the following lemma, where we denote the norm in \mathcal{C}^s by $\|\cdot\|_{\mathcal{C}^s}$.

Lemma 2.3. *Let u be the solution of (1.5). If $\Omega_0 \subset \Omega_1 \subset \Omega$ is such that Ω_1 does not contain any corner of Ω and the distance between $\partial\Omega_1 \cap \Omega$ and $\partial\Omega_0 \cap \Omega$ is positive and if the triangulations \mathcal{T}_h are quasiuniform in Ω_1 , then we have, for $\beta < s \leq 1$,*

$$\|R_h u - u\|_{\mathcal{C}(\Omega_0)} \leq C_s h^{2\beta} (\|u\|_{\mathcal{C}^{2s}(\Omega_1)} + \|\Delta u\|_{H^{-1+s}}) \leq Ch^{2\beta} (\|u\|_{\mathcal{C}^{2s}(\Omega_1)} + \|u\|_{\dot{H}^{1+s}}),$$

Optimal order $O(h^2)$ and $O(h)$ convergence in L_2 and H^1 , respectively, and almost optimal $O(h^2)$ convergence in the maximum-norm, may be obtained by systematically refining the triangulations toward the corners of Ω . Such triangulations can be defined as follows. In [8, Theorem 4.4.3.7], it is shown that any corner of Ω gives rise to singularities, expressed in terms of polar coordinates centered at the corner under consideration of the form $S_{jm}(r, \theta) = \eta_j(r) r^{\beta_{jm}} \sin(\beta_{jm}\theta)$, with $\beta_{jm} = m\pi/\omega_j \in (0, 2)$, where ω_j is the interior angle, m is a positive integer, and η_j a cutoff-function. For the reentrant corner we may have $m = 1$ or 2 , for convex corners with $\omega_j \in (\frac{1}{2}\pi, \pi)$ only $m = 1$ is possible, and for $\omega_j \leq \frac{1}{2}\pi$ no such singularity occurs.

Let $d(x)$ denote the distance to a corner, where a singularity, as described previously, occurs and, with $d_j = 2^{-j}$, let

$$\Omega_j = \{x \in \Omega : d_j/2 \leq d(x) \leq d_j\}, \quad \text{for } j = 0, \dots, J.$$

Furthermore, let $\Omega'_j = \Omega_{j-1} \cup \Omega_j \cup \Omega_{j+1}$ and $\Omega_I = \{x \in \Omega : d(x) \leq d_J/2\}$. We also assume that the mesh is locally quasiuniform on each Ω'_j . For each of the corners, β denotes the minimal corresponding β_{jm} . We now choose J such that $d_J \approx h^{2/\beta}$, where h denotes the mesh size in the interior of the domain, and $\gamma \geq 2/\beta$ such that, with ϵ any small positive number,

$$(i) \quad h_j \leq Ch d_j^{1-\beta/2+\epsilon} \quad \text{and} \quad ch^\gamma \leq h_I \leq Ch^{2/\beta}, \quad \text{with } c > 0,$$

where h_j denotes the maximal mesh size on Ω_j . We also assume that $\dim S_h \leq Ch^{-2}$ and $h_{\min} \geq h^\gamma$.

Lemma 2.4. *Let u be the solution of (1.5), if the triangulations \mathcal{T}_h satisfy (i), then we have*

$$(2.4) \quad \|R_h u - u\| + h \|\nabla(R_h u - u)\| \leq Ch^2 \|\Delta u\| = Ch^2 \|f\|.$$

Further, for any $s \in [1, 2)$ and $p < \infty$ sufficiently large, we have with $C = C_{s,p}$,

$$(2.5) \quad \|R_h u - u\|_C \leq Ch^s \|\Delta u\|_{L_p} = Ch^s \|f\|_{L_p}.$$

We note that assumption (i) corresponds to assumption (ii) of [5], and that (2.4) is true also for a weaker assumption on the triangulations. However, here we shall only employ (2.4) to show a maximum-norm estimate that also requires (2.5).

We turn now to the parabolic problem (1.1). A basic weak solution of (1.1) is such that $u \in L_2(0, T; H_0^1)$, with $u_t \in L_2(0, T; H^{-1})$ for any $T > 0$, and a unique such solution exists if $v \in L_2$ and $f \in L_2(0, T; H^{-1})$. However, in our search for maximal order convergence, the following stronger regularity results, expressed in terms of the data v and f , will be needed. We will use the notation

$$(2.6) \quad g_0 = u_t(0) = \Delta v + f(0), \text{ for } v \in \dot{H}^2, f(0) \in L_2.$$

Note that $v \in \dot{H}^2$ contains the compatibility condition $v = 0$ on $\partial\Omega$ between initial data and the boundary condition in (1.1).

Lemma 2.5. *Let u be the solution of (1.1) and assume $v = 0$ on $\partial\Omega$. Then we have, for $t \leq T$,*

$$(2.7) \quad \int_0^t \|u_t\|_{H^1}^2 d\tau \leq C \left(\|g_0\|^2 + \int_0^t \|f_t\|_{H^{-1}}^2 d\tau \right),$$

and if in addition $g_0 = 0$ on $\partial\Omega$ then

$$(2.8) \quad \int_0^t (\|\Delta u_t\|^2 + \|u_{tt}\|^2) d\tau \leq C \left(\|g_0\|_{H^1}^2 + \int_0^t \|f_t\|^2 d\tau \right).$$

Further, for $0 \leq s < 1$, with $C = C_s$,

$$(2.9) \quad \int_0^t (\|u_t\|_{\dot{H}^{1+s}} + \|u_{tt}\|_{\dot{H}^{-1+s}}) d\tau \leq C \left(\|g_0\| + \int_0^t \|f_t\| d\tau \right),$$

and if $g_0 \in H^\epsilon$ for $0 < \epsilon < \frac{1}{2}$, with $C = C_\epsilon$

$$(2.10) \quad \int_0^t (\|u_t\|_{\dot{H}^2} + \|u_{tt}\|) d\tau \leq C \left(\|g_0\|_{H^\epsilon} + \int_0^t \|f_t\|_{H^\epsilon} d\tau \right).$$

For comparison with the finite volume results to be shown in Section 3 we state some error estimates for the spatially semidiscrete finite element approximation (1.8) of the solution of (1.1).

Theorem 2.1. *Let \underline{u}_h and u be the solutions of (1.8) and (1.1) with $v = 0$ on $\partial\Omega$ and let g_0 be defined by (2.6). Then if $v_h = R_h v$, we have, for $t \leq T$,*

$$\|\underline{u}_h(t) - u(t)\| \leq Ch^{2\beta} \left(\|\Delta v\| + \|g_0\| + \int_0^t \|f_t\| d\tau \right)$$

and

$$\|\nabla \underline{u}_h(t) - \nabla u(t)\| \leq Ch^\beta \left(\|\Delta v\| + \|g_0\| + \int_0^t \|f_t\| d\tau + \left(\int_0^t \|f_t\|_{H^{-1}}^2 d\tau \right)^{1/2} \right).$$

Also we have the following maximum-norm error estimates.

Theorem 2.2. *Under the assumptions of Theorem 2.1, if the triangulations \mathcal{T}_h are such that $h_{\min} \geq Ch^\gamma$ for some $\gamma > 0$, we have, for $t \leq T$ and $s \in (0, \beta)$,*

$$\|\underline{u}_h(t) - u(t)\|_C \leq C_s h^s \left(\|v\|_{C^\beta} + \|f(0)\| + \int_0^t \|f_t\| d\tau + \left(\int_0^t \|f_t\|_{H^{-1}}^2 d\tau \right)^{1/2} \right).$$

Further if $\Omega_0 \subset \Omega_1 \subset \Omega$ is such that Ω_1 does not contain any corner of Ω and the distance between $\partial\Omega_1 \cap \Omega$ and $\partial\Omega_0 \cap \Omega$ is positive and if the triangulations \mathcal{T}_h are quasiuniform in Ω_1 and $g_0 = 0$ on $\partial\Omega$, then we have, for $s \in (\beta, 1)$ and $t \leq T$,

$$\|\underline{u}_h(t) - u(t)\|_{C(\Omega_0)} \leq Ch^{2\beta} \ell_h^{1/2} \left(\|u(t)\|_{C^{2s}(\Omega_1)} + \|\Delta v\| + \|g_0\|_{H^1} + \left(\int_0^t \|f_t\|^2 d\tau \right)^{1/2} \right).$$

Note that the first term in the parenthesis is finite provided v and f are smooth in the interior of Ω . In the presence of the appropriate refinements the convergence is almost $O(h^2)$.

Theorem 2.3. *Under the assumptions of Theorem 2.1, if the triangulations \mathcal{T}_h satisfy (i) and $g_0 = 0$ on $\partial\Omega$, then we have, for any $s \in [0, 2)$ and $t \leq T$,*

$$\|\underline{u}_h(t) - u(t)\|_C \leq C_s h^s \left(\|g_0\|_{H^1} + \|f(0)\|_C + \int_0^t \|f_t\|_C d\tau + \left(\int_0^t \|f_t\|^2 d\tau \right)^{1/2} \right).$$

3. THE SEMIDISCRETE FINITE VOLUME METHOD FOR THE PARABOLIC PROBLEM

We begin this section by recalling some basic material concerning the finite volume method, cf. [1, 2, 6, 7, 9], and then proceed with our error bounds.

We shall first rewrite the Petrov-Galerkin method (1.3) as a Galerkin method. For this purpose, we introduce the interpolation operator $J_h : \mathcal{C} \mapsto Y_h$ by

$$J_h u(x) = \sum_{z \in Z_h^0} u(z) \Psi_z(x),$$

where the set $\{\Psi_z : z \in Z_h^0\}$, with Ψ_z the characteristic function of the finite volume V_z , is a basis of Y_h . We recall that the bilinear form $(\chi, J_h \psi)$ is symmetric, positive definite on S_h , thus an inner product, and that the corresponding discrete norm is equivalent to the L_2 norm, uniformly in h , i.e., with $C \geq c > 0$,

$$(3.1) \quad c \|\chi\| \leq \|\|\chi\|\| \leq C \|\chi\|, \quad \forall \chi \in S_h, \quad \text{where } \|\|\chi\|\| \equiv (\chi, J_h \chi)^{1/2}.$$

It is well-known, cf., e.g., [1], that

$$(3.2) \quad a(\chi, \psi) = a_h(\chi, J_h \psi), \quad \forall \chi, \psi \in S_h.$$

It follows then that there exists $c > 0$, such that

$$(3.3) \quad a_h(\chi, J_h \chi) \geq c \|\nabla \chi\|^2, \quad \forall \chi \in S_h.$$

With this notation, (1.3) may equivalently be written in Galerkin form as

$$(u_{h,t}, J_h \chi) + a_h(u_h, J_h \chi) = (f, J_h \chi), \quad \forall \chi \in S_h, \quad t > 0, \quad \text{with } u_h(0) = v_h.$$

In the analysis we shall need the error functional ε_h , defined by

$$\varepsilon_h(f, \chi) = (f, J_h \chi) - (f, \chi), \quad \forall f \in H^s, \quad -\frac{1}{2} < s \leq 1, \quad \chi \in S_h,$$

and recall the following bound, cf. [2, Lemma 5.1]:

Lemma 3.1. *Let $f \in H^s$, with $s \in [0, 1]$. Then we have*

$$|\varepsilon_h(f, \chi)| \leq Ch^{i+s} \|f\|_{H^s} \|\chi\|_{H^i}, \quad \forall \chi \in S_h, \quad i = 0, 1.$$

Our next purpose is to derive an L_2 norm error estimate for the semidiscrete finite volume method (1.3), using the finite volume elliptic projection. The proof is based on the following error bound for the latter. We note that (3.5) requires more regularity than the corresponding result for the finite element projection in (2.2).

Lemma 3.2. *Let u be the solution of (1.5). Then we have, with $C = C_s$,*

$$(3.4) \quad \|\nabla(\tilde{R}_h u - u)\| \leq Ch^\beta \|\Delta u\|_{H^{-1+s}} \leq Ch^\beta \|u\|_{\dot{H}^{1+s}}, \quad \text{for } \beta < s \leq 1.$$

Further

$$(3.5) \quad \|\tilde{R}_h u - u\| \leq Ch^{2\beta} \|\Delta u\|_{H^{2\beta-1}}.$$

Proof. The estimate (3.4) is shown in [3, Theorem 5.2]. For the proof of (3.5) we employ a duality argument. For $\psi \in H_0^1$ satisfying $-\Delta\psi = \tilde{R}_h u - u$, we have

$$\|\tilde{R}_h u - u\|^2 = a(\tilde{R}_h u - u, \psi - R_h\psi) + a(\tilde{R}_h u - u, R_h\psi) = I + II.$$

For the first term we obtain, using (3.4) and (2.2), for $s = 1$ and the fact that $2\beta - 1 > 0$,

$$\begin{aligned} |I| &\leq \|\nabla(\tilde{R}_h u - u)\| \|\nabla(R_h\psi - \psi)\| \leq Ch^{2\beta} \|\Delta u\| \|\Delta\psi\| \\ &\leq Ch^{2\beta} \|\Delta u\|_{H^{2\beta-1}} \|\tilde{R}_h u - u\|. \end{aligned}$$

To bound now the second term we note that by (3.2), (1.11) and (1.4), we get

$$a(\tilde{R}_h u, R_h\psi) = a_h(\tilde{R}_h u, J_h R_h\psi) = a_h(u, J_h R_h\psi) = -(\Delta u, J_h R_h\psi),$$

so that $II = \varepsilon_h(\Delta u, R_h\psi)$, and hence, by Lemma 3.1,

$$\begin{aligned} |II| &\leq Ch^{2\beta} \|\Delta u\|_{H^{2\beta-1}} \|\nabla R_h\psi\| \leq Ch^{2\beta} \|\Delta u\|_{H^{2\beta-1}} \|\nabla\psi\| \\ &\leq Ch^{2\beta} \|\Delta u\|_{H^{2\beta-1}} \|\tilde{R}_h u - u\|. \end{aligned}$$

Together these estimates show

$$\|\tilde{R}_h u - u\|^2 \leq Ch^{2\beta} \|\Delta u\|_{H^{2\beta-1}} \|\tilde{R}_h u - u\|,$$

which completes the proof. \square

We now show an L_2 norm error estimate for (1.3) using the finite volume projection \tilde{R}_h . Here and below we denote

$$(3.6) \quad g_1 = u_{tt}(0) = \Delta g_0 + f_t(0) \text{ for } g_0 \in \dot{H}^2, \quad f_t(0) \in L_2.$$

Theorem 3.1. *Let u_h and u be the solutions of (1.3) and (1.1), respectively, and assume $v = g_0 = 0$ on $\partial\Omega$. Then, if $v_h = \tilde{R}_h v$, we have, for $t \leq T$,*

$$\|u_h(t) - u(t)\| \leq Ch^{2\beta} \left(\|\Delta v\|_{H^1} + \|g_1\| + \int_0^t (\|f_{tt}\| + \|f_t\|_{H^1}) d\tau \right).$$

Proof. Writing $u_h - u = \tilde{\vartheta} + \tilde{\varrho}$ as in (1.12) we find by (3.5)

$$(3.7) \quad \begin{aligned} \|\tilde{\varrho}(t)\| &\leq \|\tilde{\varrho}(0)\| + \int_0^t \|\tilde{\varrho}_t\| d\tau \leq Ch^{2\beta} \left(\|\Delta v\|_{H^{2\beta-1}} + \int_0^t \|\Delta u_t\|_{H^{2\beta-1}} d\tau \right) \\ &\leq Ch^{2\beta} \left(\|\Delta v\|_{H^1} + \int_0^t (\|u_{tt}\|_{H^1} + \|f_t\|_{H^1}) d\tau \right), \end{aligned}$$

where in the last step we have used the fact that $2\beta - 1 \in (0, 1)$ and $\Delta u_t = u_{tt} - f_t$. Since u_t satisfies (1.1), with f and v replaced by f_t and $u_t(0) = g_0$, respectively, the regularity estimate (2.9) shows

$$(3.8) \quad \int_0^t \|u_{tt}\|_{H^1} \leq C \left(\|g_1\| + \int_0^t \|f_{tt}\| \, d\tau \right),$$

which applied to (3.7) bounds $\tilde{\varrho}$ as desired. We now turn to $\tilde{\vartheta}$, which satisfies (1.13). Thus choosing $\eta = J_h \tilde{\vartheta}$ we get

$$(\tilde{\vartheta}_t, J_h \tilde{\vartheta}) + a_h(\tilde{\vartheta}, J_h \tilde{\vartheta}) = -(\tilde{\varrho}_t, J_h \tilde{\vartheta}),$$

and hence by standard energy arguments we obtain

$$\|\tilde{\vartheta}(t)\| \leq C \int_0^t \|\tilde{\varrho}_t\| \, d\tau.$$

Finally, in view of (3.7) and (3.8), this completes the proof. \square

We now show an L_2 norm error estimate for (1.3) using instead the finite element Ritz projection R_h in the analysis. Note that in this case the regularity requirements on data are weaker than in Theorem 3.1.

Theorem 3.2. *Let u_h and u be the solutions of (1.3) and (1.1), respectively, and assume $v = 0$ on $\partial\Omega$. Then, if $v_h = R_h v$, we have, for $t \leq T$,*

$$\|u_h(t) - u(t)\| \leq Ch^{2\beta} \left(\|\Delta v\| + \|g_0\| + \left(\int_0^t (\|f_t\|^2 + \|f\|_{H^1}^2) \, d\tau \right)^{1/2} \right).$$

Proof. We write $u_h - u = \vartheta + \varrho$, as in (1.14). Then (2.2), the fact that $\|v\|_{\dot{H}^{1+s}} \leq C\|v\|_{\dot{H}^2} = C\|\Delta v\|$, and (2.9) give for $s = (1 + \beta)/2$,

$$(3.9) \quad \begin{aligned} \|\varrho(t)\| &\leq \|\varrho(0)\| + \int_0^t \|\varrho_t\| \, d\tau \leq Ch^{2\beta} \left(\|v\|_{\dot{H}^{1+s}} + \int_0^t \|u_t\|_{\dot{H}^{1+s}} \, d\tau \right) \\ &\leq Ch^{2\beta} \left(\|\Delta v\| + \|g_0\| + \int_0^t \|f_t\| \, d\tau \right), \end{aligned}$$

which yields the desired estimate for ϱ .

We turn to the estimation of ϑ , which satisfies the equation (1.15). In view of (3.2), (1.4) and (1.9), we get

$$(3.10) \quad a_h(\varrho, J_h \chi) = a(R_h u, \chi) + (\Delta u, J_h \chi) = \varepsilon_h(\Delta u, \chi), \quad \forall \chi \in S_h.$$

Using this, (1.15) with $\eta = J_h \vartheta$ is transformed into

$$(\vartheta_t, J_h \vartheta) + a_h(\vartheta, J_h \vartheta) = -(\varrho_t, J_h \vartheta) - \varepsilon_h(\Delta u, \vartheta).$$

In view of Lemma 3.1 and the symmetry of $(\chi, J_h \psi)$ on S_h , this shows

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\vartheta\|^2 + a_h(\vartheta, J_h \vartheta) &\leq C \|\varrho_t\| \|\vartheta\| + Ch^{2\beta} \|\Delta u\|_{H^{2\beta-1}} \|\vartheta\|_{H^1} \\ &\leq C \|\varrho_t\| \|\vartheta\| + Ch^{2\beta} \left(\|u_t\|_{H^1} + \|f\|_{H^1} \right) \|\nabla \vartheta\|. \end{aligned}$$

Using (3.3) to kick back $\|\nabla\vartheta\|$, and integrating over $[0, t]$, we obtain, in view of (3.1), (2.7), and the fact that $\vartheta(0) = 0$,

$$\begin{aligned}\|\vartheta(t)\|^2 &\leq C \int_0^t \|\varrho_t\| \|\vartheta\| d\tau + Ch^{4\beta} \int_0^t (\|u_t\|_{H^1}^2 + \|f\|_{H^1}^2) d\tau \\ &\leq C \int_0^t \|\varrho_t\| \|\vartheta\| d\tau + Ch^{4\beta} \left(\|g_0\|^2 + \int_0^t (\|f_t\|_{H^{-1}}^2 + \|f\|_{H^1}^2) d\tau \right).\end{aligned}$$

Setting $\Theta(t) \equiv \sup_{0 < s \leq t} \|\vartheta(s)\|$, this shows

$$\|\vartheta(t)\|^2 \leq \Theta(t)^2 \leq C \left(\int_0^t \|\varrho_t\| d\tau \right) \Theta(t) + Ch^{4\beta} \left(\|g_0\|^2 + \int_0^t (\|f_t\|_{H^{-1}}^2 + \|f\|_{H^{2\beta-1}}^2) d\tau \right),$$

which, in view of (3.9), gives the desired bound for ϑ . \square

Next, we show an $O(h^\beta)$ estimate for the gradient of the error.

Theorem 3.3. *Under the assumptions of Theorem 3.2, we have, for $t \leq T$,*

$$\|\nabla(u_h(t) - u(t))\| \leq Ch^\beta \left(\|\Delta v\| + \|g_0\| + \left(\int_0^t \|f_t\|^2 d\tau \right)^{1/2} \right).$$

Proof. In view of (2.2) we have, for $s = (1 + \beta)/2$,

$$(3.11) \quad \|\nabla\varrho(t)\| \leq \|\nabla\varrho(0)\| + \int_0^t \|\nabla\varrho_t\| d\tau \leq Ch^\beta \left(\|v\|_{\dot{H}^{1+s}} + \int_0^t \|u_t\|_{\dot{H}^{1+s}} d\tau \right),$$

and (2.9) then gives the desired bound for ϱ . To bound ϑ we now choose $\eta = J_h\vartheta_t$ in (1.15), and using (3.10) with the fact that $a_h(\vartheta, J_h\vartheta_t) = \frac{1}{2} \frac{d}{dt} a(\vartheta, \vartheta)$, we get

$$(3.12) \quad \|\vartheta_t\|^2 + \frac{1}{2} \frac{d}{dt} a(\vartheta, \vartheta) = -(\varrho_t, J_h\vartheta_t) - \varepsilon_h(\Delta u, \vartheta_t).$$

Substituting $-\Delta u = f - u_t$ yields

$$\|\vartheta_t\|^2 + \frac{1}{2} \frac{d}{dt} a(\vartheta, \vartheta) = -(\varrho_t, J_h\vartheta_t) - \varepsilon_h(u_t, \vartheta_t) + \frac{d}{dt} \varepsilon_h(f, \vartheta) - \varepsilon_h(f_t, \vartheta).$$

Integrating this relation over $[0, t]$, using $\vartheta(0) = 0$, together with (3.1) and Lemma 3.1, we find

$$\begin{aligned}\int_0^t \|\vartheta_t\|^2 d\tau + \frac{1}{2} \|\nabla\vartheta(t)\|^2 &\leq Ch \|f(t)\| \|\nabla\vartheta(t)\| \\ &\quad + \int_0^t (\|\varrho_t\| \|\vartheta_t\| + Ch(\|u_t\|_{H^1} \|\vartheta_t\| + \|f_t\| \|\nabla\vartheta\|)) d\tau.\end{aligned}$$

This together with (2.3), and the fact that $\beta < 1$ give

$$\|\nabla\vartheta(t)\|^2 \leq Ch^{2\beta} \left(\|f(t)\|^2 + \int_0^t (\|u_t\|_{H^1}^2 + \|f_t\|^2) d\tau \right) + C \int_0^t \|\nabla\vartheta\|^2 d\tau.$$

Using Gronwall's lemma, the estimate

$$\|f(t)\|^2 \leq C \left(\|\Delta v\|^2 + \|g_0\|^2 + \int_0^t \|f_t\|^2 d\tau \right),$$

and (2.7), we finally get,

$$(3.13) \quad \|\nabla\vartheta(t)\|^2 \leq Ch^{2\beta} \left(\|\Delta v\|^2 + \|g_0\|^2 + \int_0^t \|f_t\|^2 d\tau \right),$$

which gives the desired bound for ϑ . \square

Based on the above analysis of $\nabla\vartheta$ we can show the following “super”-closeness of the gradients of u_h and $R_h u$.

Lemma 3.3. *Under the assumptions of Theorem 3.2, let $g_0 = 0$ on $\partial\Omega$ and g_1 be defined by (3.6). Then we have, for $t \leq T$,*

$$\|\nabla\vartheta(t)\| \leq Ch^{2\beta} \left(\|\Delta v\|_{H^1} + \|g_1\| + \left(\int_0^t (\|f_t\|_{H^1}^2 + \|f_{tt}\|_{H^{-1}}^2) d\tau \right)^{1/2} \right).$$

Proof. Rewriting the right hand side of equation (3.12) in the form

$$(3.14) \quad \|\vartheta_t\|^2 + \frac{1}{2} \frac{d}{dt} a(\vartheta, \vartheta) = -(\varrho_t, J_h \vartheta_t) - \frac{d}{dt} \varepsilon_h(\Delta u, \vartheta) + \varepsilon_h(\Delta u_t, \vartheta),$$

integrating this over $[0, t]$, using $\vartheta(0) = 0$, together with Lemma 3.1, and (3.1), we find

$$\begin{aligned} \int_0^t \|\vartheta_t\|^2 d\tau + \frac{1}{2} \|\nabla\vartheta(t)\|^2 &\leq Ch^{2\beta} \|\Delta u(t)\|_{H^{2\beta-1}} \|\nabla\vartheta(t)\| \\ &\quad + \int_0^t (\|\varrho_t\| \|\vartheta_t\| + Ch^{2\beta} \|\Delta u_t\|_{H^{2\beta-1}} \|\nabla\vartheta\|) d\tau. \end{aligned}$$

Together with (2.2) for $s = 1$ and the fact that $2\beta - 1 \in (0, 1)$, this gives

$$\|\nabla\vartheta(t)\|^2 \leq Ch^{4\beta} \left(\|\Delta v\|_{H^1}^2 + \int_0^t (\|u_{tt}\|_{H^1}^2 + \|f_t\|_{H^1}^2) d\tau \right) + C \int_0^t \|\nabla\vartheta\|^2 d\tau.$$

Since u_t satisfies (1.1), with f and v replaced by f_t and $u_t(0) = g_0$, respectively, the regularity estimate (2.7) shows

$$(3.15) \quad \int_0^t \|u_{tt}\|_{H^1}^2 \leq C \left(\|g_1\|^2 + \int_0^t \|f_{tt}\|_{H^{-1}}^2 d\tau \right).$$

Using this together with Gronwall’s lemma, we obtain the desired estimate. \square

We now turn to error estimates in maximum-norm. If the triangulations are such that $h_{\min} \geq Ch^\gamma$, for some $\gamma > 0$ then we can get $\|\varrho\|_C = O(h^s)$, $s \in (0, \beta)$, cf. Lemma 2.2 and $\|\vartheta\|_C = O(h^\beta \ell_h^{1/2})$. In [5], for the finite element approximation, we demonstrated that the convergence rate is higher in maximum-norm over a subdomain that is away from the corners or under properly refined (near the corner) meshes \mathcal{T}_h . This applies also to the corresponding finite volume approximations.

First we show a global maximum-norm error estimate. Note that the maximum-norm error for the finite element method, cf. Theorem 2.2, is almost of order $O(h^\beta)$, under almost the same regularity assumptions.

Theorem 3.4. *Under the assumptions of Theorem 3.2, if the triangulations \mathcal{T}_h are such that $h_{\min} \geq Ch^\gamma$ for some $\gamma > 0$, then we have, for $s \in (0, \beta)$ and $t \leq T$, with $C = C_s$,*

$$\|u_h(t) - u(t)\|_C \leq Ch^s \left(\|\Delta v\| + \|g_0\| + \left(\int_0^t \|f_t\|^2 d\tau \right)^{1/2} \right).$$

Proof. We have by Lemma 2.2, with $s_1 \in (s, \beta)$,

$$\|\varrho(t)\|_C \leq Ch^s \|u(t)\|_{\dot{H}^{1+s_1}} \leq Ch^s \left(\|v\|_{\dot{H}^{1+s_1}} + \int_0^t \|u_t\|_{\dot{H}^{1+s_1}} d\tau \right),$$

which is bounded as desired by (2.9).

In [5] we showed that the following discrete Sobolev type inequality, is valid for triangulations satisfying the condition assumed in this theorem

$$(3.16) \quad \|\chi\|_c \leq C\ell_h^{1/2}\|\nabla\chi\|, \quad \forall\chi \in \mathcal{S}_h.$$

Hence, in view of (3.13) we get

$$\|\vartheta(t)\|_c \leq Ch^\beta\ell_h^{1/2}\left(\|\Delta v\| + \|g_0\| + \left(\int_0^t \|f_t\|^2 d\tau\right)^{1/2}\right),$$

which shows the desired estimate for ϑ . \square

We remark that under the stronger assumption $\gamma = 1$, i.e., when the triangulations \mathcal{T}_h are globally quasiuniform, one can obtain an $O(h^\beta\ell_h)$ maximum-norm estimate, under marginally weaker regularity assumptions on data, as remarked after Lemma 2.2.

Next, we derive an error estimate away from the corners of Ω .

Theorem 3.5. *Under the assumptions of Theorem 3.4, let $g_0 = 0$ on $\partial\Omega$ and g_1 be defined by (3.6). If $\Omega_0 \subset \Omega_1 \subset \Omega$ is such that Ω_1 does not contain any corner of Ω and the distance between $\partial\Omega_1 \cap \Omega$ and $\partial\Omega_0 \cap \Omega$ is positive, and if the triangulations \mathcal{T}_h are quasiuniform in Ω_1 , then we have, for $t \leq T$,*

$$\begin{aligned} \|u_h(t) - u(t)\|_{c(\Omega_0)} &\leq Ch^{2\beta}\ell_h^{1/2}\left(\|\Delta v\|_{H^1} + \|g_0\| + \|g_1\| \right. \\ &\quad \left. + \left(\int_0^t (\|f_t\|_{H^1}^2 + \|f_{tt}\|_{H^{-1}}^2) d\tau\right)^{1/2}\right). \end{aligned}$$

Proof. By Lemma 2.3 we have, with $s = (1 + \beta)/2$,

$$\|\varrho(t)\|_{c(\Omega_0)} \leq Ch^{2\beta}\left(\|u(t)\|_{c^{2s}(\Omega_1)} + \|v\|_{\dot{H}^{1+s}} + \int_0^t \|u_t\|_{\dot{H}^{1+s}} d\tau\right),$$

which is bounded as desired in view of the estimate

$$\begin{aligned} \|u(t)\|_{c^{2s}(\Omega_1)} &\leq C\|\Delta u(t)\|_{H^1} \leq C\left(\|\Delta v\|_{H^1} + \int_0^t \|\Delta u_t\|_{H^1} d\tau\right) \\ &\leq C\left(\|\Delta v\|_{H^1} + \int_0^t (\|u_{tt}\|_{H^1} + \|f_t\|_{H^1}) d\tau\right) \end{aligned}$$

the regularity estimate (3.8), the fact that $\|v\|_{\dot{H}^{1+s}} \leq C\|\Delta v\|$ and (2.9). Further, using the supercloseness result of Lemma 3.3 together with (3.16) we bound ϑ as desired, and thus complete the proof. \square

We also have the following result showing almost $O(h^2)$ convergence in the presence of appropriate refinements.

Theorem 3.6. *Under the assumptions of Lemma 3.3, if the triangulations \mathcal{T}_h be refined as in Lemma 2.4, then we have, for $t \leq T$, with $C = C_s$ and $s \in (0, 2)$,*

$$\|u_h(t) - u(t)\|_c \leq Ch^s\left(\|\Delta v\|_{H^1} + \|g_1\| + \left(\int_0^t (\|f_t\|_{H^1}^2 + \|f_{tt}\|^2) d\tau\right)^{1/2}\right).$$

Proof. To bound ϱ we use (2.5) for p sufficiently large and a standard Sobolev inequality, to obtain with $C = C_{s,p}$,

$$\begin{aligned} \|\varrho(t)\|_C &\leq Ch^s \|\Delta u(t)\|_{L_p} \leq Ch^s \|\Delta u(t)\|_{H^1} \\ &\leq Ch^s \left(\|\Delta v\|_{H^1} + \int_0^t (\|u_{tt}\|_{H^1} + \|f_t\|_{H^1}) d\tau \right), \end{aligned}$$

which in view (3.8) gives the desired bound for ϱ .

We next derive a superconvergent $O(h^2)$ order estimate for $\|\nabla\vartheta\|$ based on the L_2 norm error bound of (2.4). For this we follow the proof of Lemma 3.3, stating now (3.14) and obtain this time after the application of Gronwall's lemma

$$\begin{aligned} \|\nabla\vartheta(t)\|^2 &\leq Ch^4 \left(\|\Delta v\|_{H^1}^2 + \int_0^t \|\Delta u_t\|_{H^1}^2 d\tau \right) \\ &\leq Ch^4 \left(\|\Delta v\|_{H^1}^2 + \int_0^t (\|u_{tt}\|_{H^1}^2 + \|f_t\|_{H^1}^2) d\tau \right), \end{aligned}$$

which, in view of the regularity estimate (3.15) and (3.16) completes the proof. \square

We note that under less stringent assumptions on the triangulations \mathcal{T}_h than (i), one can show optimal order $O(h^2)$ and $O(h)$ convergence, in L_2 and H^1 norm, respectively, for the error $u_h - u$, cf. [5] for a corresponding result for the finite element approximation \underline{u}_h

4. THE BACKWARD EULER FULLY DISCRETE SCHEME

In [5], in addition to the semidiscrete finite element problem (1.8), also fully discrete methods were considered. These were obtained by discretizing (1.8) in time by the backward Euler and Crank-Nicolson methods. The time discretization resulted in slightly higher regularity requirements on data than those summarized in Section 2 above and we refer to [5] for details.

In this section, by application of our analysis of the semidiscrete finite volume problem (1.3) to a fully discrete scheme, we will show some error estimates for the discretization in time by the *Backward Euler* method. Letting k denote the time step, U^n the approximation in S_h of $u(t)$ at $t = t_n = nk$, and $\bar{\partial}U^n = (U^n - U^{n-1})/k$, we consider the fully discrete scheme

$$(4.1) \quad (\bar{\partial}U^n, \eta) + a_h(U^n, \eta) = (f(t_n), \eta), \quad \forall \eta \in Y_h, \quad \text{with } U^0 = v_h = R_h v.$$

We first show the following error estimate in L_2 , with g_0 defined in (2.6).

Theorem 4.1. *Let U^n and $u(t_n)$ be the solutions of (4.1) and (1.1), respectively, and assume $v = 0$ on $\partial\Omega$. Then we have, for $t_n \leq T$ and $\epsilon \in (0, \frac{1}{2})$, with $C = C_\epsilon$,*

$$\|U^n - u(t_n)\| \leq C(h^{2\beta} + k) \left(\|\Delta v\| + \|g_0\|_{H^\epsilon} + \left(\int_0^{t_n} (\|f\|_{H^1}^2 + \|f_t\|_{H^1}^2) d\tau \right)^{1/2} \right).$$

Proof. Analogously to (1.14) we write

$$U^n - u(t_n) = (U^n - R_h u(t_n)) + (R_h u(t_n) - u(t_n)) = \vartheta^n + \varrho^n.$$

Here ϱ^n is bounded as desired by (3.9). To bound ϑ^n we note that

$$(4.2) \quad \begin{aligned} (\bar{\partial}\vartheta^n, \eta) + a_h(\vartheta^n, \eta) &= -(\bar{\partial}\varrho^n, \eta) + (u_t^n - \bar{\partial}u^n, \eta) - a_h(\varrho^n, \eta) \\ &= -(\omega^n, \eta) - a_h(\varrho^n, \eta), \quad \forall \eta \in Y_h, \end{aligned}$$

where

$$(4.3) \quad \omega^n = \omega_1^n + \omega_2^n = (R_h - I)\bar{\partial}u(t_n) + (\bar{\partial}u(t_n) - u_t(t_n)).$$

Choosing $\eta = J_h \vartheta^n$ in (4.2) we obtain, in view of (3.10),

$$\begin{aligned} & \frac{1}{2k} (\|\vartheta^n\|^2 - \|\vartheta^{n-1}\|^2) + \frac{1}{2k} \|\vartheta^n - \vartheta^{n-1}\|^2 + a_h(\vartheta^n, J_h \vartheta^n) \\ &= -(\omega^n, J_h \vartheta^n) - \varepsilon_h(u_t^n - f^n, \vartheta^n) \\ &= -(\omega_1^n, J_h \vartheta^n) - (\omega_2^n, \vartheta^n) - \varepsilon_h(\bar{\partial}u^n - f^n, \vartheta^n). \end{aligned}$$

Multiplying this by $2k$, employing Lemma 3.1 and (3.1), and kicking back $\|\nabla \vartheta^n\|$, we get

$$\|\vartheta^n\|^2 \leq \|\vartheta^{n-1}\|^2 + Ck(\|\omega_1^n\| + \|\omega_2^n\|)\|\vartheta^n\| + Ckh^4(\|\bar{\partial}u^n\|_{H^1}^2 + \|f^n\|_{H^1}^2).$$

Let now $\Theta^n = \max_{0 \leq j \leq n} \|\vartheta^j\|$. Then, since $\vartheta^0 = 0$ and using again (3.1),

$$(4.4) \quad \|\vartheta^n\|^2 \leq (\Theta^n)^2 \leq Ck \sum_{j=1}^n (\|\omega_1^j\| + \|\omega_2^j\|)\Theta^n + Ckh^4 \sum_{j=1}^n (\|\bar{\partial}u^j\|_{H^1}^2 + \|f^j\|_{H^1}^2).$$

To bound the first term in the right hand side of (4.4), we employ the inequality

$$(4.5) \quad k \sum_{j=1}^n |\bar{\partial}g^j|^p \leq Ck \sum_{j=1}^n (k^{-1} \int_{t_{j-1}}^{t_j} |g_t| d\tau)^p \leq C \int_0^{t_n} |g_t|^p d\tau, \quad 1 \leq p < +\infty,$$

where $(X, |\cdot|)$ is a normed linear space, for $p = 1$, (2.2), with $\beta < s < 1$,

$$(4.6) \quad \begin{aligned} Ck \sum_{j=1}^n (\|\omega_1^j\| + \|\omega_2^j\|) &\leq Ck \sum_{j=1}^n (h^{2\beta} \|\bar{\partial}u^j\|_{\dot{H}^{1+s}} + \int_{t_{j-1}}^{t_j} \|u_{tt}\| d\tau) \\ &\leq Ch^{2\beta} \int_0^{t_n} \|u_t\|_{\dot{H}^{1+s}} d\tau + Ck \int_0^{t_n} \|u_{tt}\| d\tau. \end{aligned}$$

To bound now the last term in (4.4), we use the inequality

$$k \sum_{j=1}^n |g^j|^2 \leq C \int_0^{t_n} (|g|^2 + k^2 |g_t|^2) d\tau \leq C \int_0^{t_n} (|g|^2 + |g_t|^2) d\tau$$

and (4.5) for $p = 2$ to obtain

$$(4.7) \quad k \sum_{j=1}^n (\|\bar{\partial}u^j\|_{H^1}^2 + \|f^j\|_{H^1}^2) \leq C \int_0^{t_n} (\|u_t\|_{H^1}^2 + \|f\|_{H^1}^2 + \|f_t\|_{H^1}^2) d\tau.$$

Finally, (4.4), (4.6) and (4.7) give

$$\begin{aligned} \|\vartheta^n\|^2 &\leq Ch^{4\beta} \left(\int_0^{t_n} \|u_t\|_{\dot{H}^{1+s}} d\tau \right)^2 + \int_0^{t_n} (\|u_t\|_{H^1}^2 + \|f\|_{H^1}^2 + \|f_t\|_{H^1}^2) d\tau \\ &\quad + Ck^2 \left(\int_0^{t_n} \|u_{tt}\| d\tau \right)^2. \end{aligned}$$

In view of (2.7) and (2.10) this completes the proof. \square

Next, we will show the following error estimate for the gradient.

Theorem 4.2. *Under the assumptions of Theorem 4.1, let $g_0 = 0$ on $\partial\Omega$. Then we have, for $t_n \leq T$,*

$$\|\nabla(U^n - u(t_n))\| \leq C(h^\beta + k) \left(\|\Delta v\| + \|g_0\|_{H^1} + \left(\int_0^{t_n} \|f_t\|^2 d\tau \right)^{1/2} \right).$$

Proof. Here ϱ^n is bounded as desired by (3.11) and (2.7). To estimate ϑ^n , we choose $\eta = J_h \bar{\partial} \vartheta^n$ in (4.2). Using (3.2) together with the identity $2a(\vartheta^n, \bar{\partial} \vartheta^n) = \bar{\partial} \|\nabla \vartheta^n\|^2 + k \|\nabla \bar{\partial} \vartheta^n\|^2$, and (3.10), we then obtain

$$\|\bar{\partial} \vartheta^n\|^2 + \frac{1}{2} (\bar{\partial} \|\nabla \vartheta^n\|^2 + k \|\nabla \bar{\partial} \vartheta^n\|^2) = -(\omega^n, J_h \bar{\partial} \vartheta^n) - \varepsilon_h(\Delta u^n, \bar{\partial} \vartheta^n).$$

Multiplying this by $2k$, using (3.1), eliminating $\|\bar{\partial} \vartheta^n\|^2$, and summing in time, we have

$$(4.8) \quad \|\nabla \vartheta^n\|^2 \leq Ck \sum_{j=1}^n \|\omega^j\|^2 + k \sum_{j=1}^n \varepsilon_h(\Delta u^j, \bar{\partial} \vartheta^j).$$

Since $\vartheta^0 = 0$, the last term can be rewritten as

$$k \sum_{j=1}^n \varepsilon_h(\Delta u^j, \bar{\partial} \vartheta^j) = \varepsilon_h(\Delta u^n, \vartheta^n) - k \sum_{j=1}^n \varepsilon_h(\bar{\partial} \Delta u^j, \vartheta^{j-1}).$$

Thus, employing this identity and Lemma 3.1 in (4.8), and using the discrete version of Gronwall's lemma, we get

$$(4.9) \quad \|\nabla \vartheta^n\|^2 \leq Ck \sum_{j=1}^n (\|\omega_1^j\|^2 + \|\omega_2^j\|^2) + Ch^2 (\|\Delta u^n\|^2 + k \sum_{j=1}^n \|\bar{\partial} \Delta u^j\|^2).$$

Using (4.5) with $p = 2$, we easily find

$$\|\Delta u^n\|^2 + k \sum_{j=1}^n \|\bar{\partial} \Delta u^j\|^2 \leq C(\|\Delta v\|^2 + \int_0^{t_n} \|\Delta u_t\|^2 d\tau).$$

Hence by (2.8) the last term in (4.9) is bounded as desired. Using (2.3) and again (4.5) with $p = 2$, we obtain

$$\begin{aligned} k \sum_{j=1}^n (\|\omega_1^j\|^2 + \|\omega_2^j\|^2) &\leq Ck \sum_{j=1}^n \left(h^{2\beta} \|\bar{\partial} u^j\|_{H^1}^2 + \left(\int_{t_{j-1}}^{t_j} \|u_{tt}\| d\tau \right)^2 \right) \\ &\leq Ch^{2\beta} \int_0^{t_n} \|u_t\|_{H^1}^2 d\tau + Ck^2 \int_0^{t_n} \|u_{tt}\|^2 d\tau. \end{aligned}$$

Then we complete the proof by applying (2.7) and (2.8) and thus estimating the remaining term of the right hand side of (4.9) as desired, i.e.,

$$(4.10) \quad \|\nabla \vartheta^n\| \leq C(h^\beta + k) \left(\|\Delta v\| + \|g_0\|_{H^1} + \left(\int_0^{t_n} \|f_t\|^2 d\tau \right)^{1/2} \right). \quad \square$$

We finally demonstrate the following time discrete version of the maximum-norm error estimate of Theorem 3.4.

Theorem 4.3. *Under the assumptions of Theorem 4.2, if the triangulations \mathcal{T}_h are such that $h_{\min} \geq Ch^\gamma$ for some $\gamma > 0$, then we have, for $0 \leq s < \beta$ and $t_n \leq T$, with $C = C_s$,*

$$\|U^n - u(t_n)\|_C \leq C(h^s + k\ell_h^{1/2})\left(\|\Delta v\| + \|g_0\|_{H^1} + \left(\int_0^{t_n} \|f_t\|^2 d\tau\right)^{1/2}\right).$$

Proof. The term ϱ^n is bounded as stated by following the proof of Theorem 3.4. The proof is completed by bounding ϑ^n , using (3.16) and (4.10), as

$$\|\vartheta^n\|_C \leq \ell_h^{1/2} \|\nabla \vartheta^n\| \leq C(h^\beta + k)\ell_h^{1/2}\left(\|\Delta v\| + \|g_0\|_{H^1} + \left(\int_0^{t_n} \|f_t\|^2 d\tau\right)^{1/2}\right). \quad \square$$

We refrain from stating and proving the straight-forward analogous results of Theorems 3.5 and 3.6.

REFERENCES

- [1] R. E. Bank and D. J. Rose. Some error estimates for the box method, *SIAM J. Numer. Anal.*, 24(4):777–787, 1987.
- [2] P. Chatzipantelidis, Finite volume methods for elliptic PDE’s: a new approach, *M2AN Math. Model. Numer. Anal.*, 36(2):307–324, 2002.
- [3] P. Chatzipantelidis and R. D. Lazarov, Error estimates for a finite volume element method for elliptic pde’s in nonconvex polygonal domains, *SIAM J. Numer. Anal.*, 42:1932–1958, 2005.
- [4] P. Chatzipantelidis, R. D. Lazarov, and V. Thomée, Error estimates for a finite volume element method for parabolic equations in convex polygonal domains, *Numer. Methods Partial Differential Equations*, 20(5):650–674, 2004.
- [5] P. Chatzipantelidis, R. D. Lazarov, V. Thomée and L. B. Wahlbin, Parabolic finite element equations in nonconvex polygonal domains, *BIT Numerical Mathematics*, 46 (suppl. 5): S113–S143, 2006.
- [6] S.-H. Chou and Q. Li, Error estimates in L^2 , H^1 and L^∞ in covolume methods for elliptic and parabolic problems: a unified approach, *Math. Comp.*, 69(229):103–120, 2000.
- [7] R. Eymard, T. Gallouët, and R. Herbin, *Finite volume methods*, In *Handbook of numerical analysis, Vol. VII*, Handbook of Numerical Analysis, VII, pages 713–1020. North-Holland, Amsterdam, 2000.
- [8] P. Grisvard, *Elliptic Problems in Nonsmooth Domains*, Pitman, Massachusetts, 1985.
- [9] R. Li, Z. Chen, and W. Wu, *Generalized difference methods for differential equations: Numerical analysis of finite volume methods*, *Monographs and Textbooks in Pure and Applied Mathematics, vol. 226*, Marcel Dekker Inc., New York, 2000.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CRETE, 714 09 HERAKLION, GREECE
E-mail address: chatzipa@math.uoc.gr

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843, USA
E-mail address: lazarov@math.tamu.edu

DEPARTMENT OF MATHEMATICS, CHALMERS UNIVERSITY OF TECHNOLOGY AND GÖTEBORG UNIVERSITY, SE-412 96 GÖTEBORG, SWEDEN
E-mail address: thomee@math.chalmers.se